

Echinacea (Coneflower)

30ct Plug

Container:

Premium 1 Gallon Pot

Finish:

6-8 weeks

When To Plant:

Late summer or early to late spring

72ct Plug

Container:

Premium 1 Gallon Pot

Finish:

8-10 weeks

When To Plant:

Mid to late summer or early spring to early summer

Growing Temperature:

68° F

Soil pH:

5.5-6.5

Fertility:

Apply 100-200 N ppm of a standard water soluble fertilizer every 7-14 days once plants begin growing. Echinacea DO NOT like to be over fertilized when growing in containers. This often leads to increased incidences of crown and root rots and a general decline in plant vigor. Use slightly acidic soil with an EC of 1.0-2.6 using the pour through method for best results.

For Controlled Release or Slow Release Fertilizer, see your preferred supplier for recommended rates for incorporation or top dressing, as it varies by fertilizer.

Potting and Timing:

To get the plants to bulk up before winter, increase their day length and do not allow them to flower.

Planting Level:

At crown level.

PGRs/Pinching:

Because night interruption lighting will cause plants to stretch, make your first growth regulator applications before lighting begins and apply again two weeks later. Discontinue lighting once the plants begin to grow.

B-Nine at 2500 ppm or Sumagic at 5 ppm works well to control overall height. Sumagic applied as a drench at 1 ppm also works well.

Holding Temperature:

50-60° F

EC Level:

1.5-2.0 pour through method

Vernalization:

Cold beneficial. An 11-14 week cold period will increase the number of flowers and overall quality of Echinacea.

Pests and Diseases:

Scout for whitefly, aphids, thrips and many types of caterpillars. Many of these insects carry plant viruses that can be transmitted to Echinacea. If conditions are favorable they can be susceptible to Botrytis, Rhizoctonia and Pythium. Echinacea are susceptible to Aster Yellows disease, which is caused by a phytoplasma transmitted by the six-spotted (aster) leafhopper. Affected plants generally show yellowing of the leaves, reddening, and stunting in the early stages of disease development. One of the most notable signs of Aster Yellows is deformed flowers.

Moisture:

Keep plugs moist until they are ready to transplant. Just before transplanting, make sure the plugs are well watered. Once they have been transplanted and are actively growing, it is important that they are not overwatered. Overwatering may result in plants that grow more slowly in containers and are more susceptible to disease. Plants should never be allowed to sit in excess water.

Lighting:

Echinacea are considered intermediate day plants, requiring 12 to 16 hours of day length to flower. Day lengths less than 12 or more than 16 hours will tend to keep the plants in a vegetative state.

When day lengths are short and flowers are desired, provide 4 hour night interruption.

Grower Tips:

Keys to overwintering Echinacea are:

- Be sure the plants are fully rooted and have a good size crown. Undersized plants will not overwinter well.
- Slowly reduce the level of feed going into fall as the temperatures cool.
- It is recommended to drench plants with a preventive fungicide treatment in late summer to prevent root and crown rots.
- A soil test should be performed to measure where the soluble salts levels are (EC). This should be in the 0.5-1.0 range using the pour through method of testing. Elevated salt levels will burn the roots and cause crown and root rots.
- Monitor soil moisture through the winter. Avoid overly wet conditions and be sure to avoid heavy condensation in the greenhouse.
- During exceptionally cold periods during the winter a heavy frost blanket will help provide extra protection from the cold.
- Avoid large temperature swings as this causes premature



WaltersGardens
INC

P.O. Box 137 • 1992 96th Ave. Zeeland, MI. 49464

Phone: 888-925-8377 • Fax: 800-752-1879

Email: sales@waltersgardens.com

www.WaltersGardens.com • www.PerennialResource.com

www.ProvenWinners.com